

Infinite monoids as geometric objects

Robert Gray
(joint work with Mark Kambites)

Semigroups and Applications
Novi Sad, June 2013

Groups, monoids, and geometry

Gromov - “Infinite groups as geometric objects” International Congress of Mathematicians address in Warsaw, 1984

There are two main inter-related strands in **geometric group theory**:

1. one seeks to understand groups by studying their actions on appropriate spaces, and
2. one seeks understanding from the intrinsic geometry of finitely generated groups endowed with word metrics.

Groups, monoids, and geometry

Gromov - “Infinite groups as geometric objects” International Congress of Mathematicians address in Warsaw, 1984

There are two main inter-related strands in **geometric group theory**:

1. one seeks to understand groups by studying their actions on appropriate spaces, and
2. one seeks understanding from the intrinsic geometry of finitely generated groups endowed with word metrics.

How about monoids and semigroups?

1. To what extent can we gain information about finitely generated monoids by studying their actions on geometric objects?
2. How much algebraic information about finitely generated monoids is encoded in the geometry of their Cayley graphs?

General philosophy

Algebra

Combinatorics

Groups

Graphs

Monoids / Semigroups

Digraphs

General philosophy

Algebra

Combinatorics

Geometry

Groups

Graphs

Metric spaces

Monoids / Semigroups

Digraphs

??

General philosophy

Algebra	Combinatorics	Geometry
Groups	Graphs	Metric spaces
Monoids / Semigroups	Digraphs	??

?? = directed metric spaces = semimetric spaces

Semimetric space = a set equipped with an asymmetric, partially-defined distance function.

Cayley graphs and the notion of quasi-isometry

G - group, generated by a finite set $A \subseteq G$,

Assume $1 \notin A$, and $A = A^{-1}$

G gives rise to a metric space (G, d_A) with word metric d_A .

Points: G


Distance: $d_A(g, h)$ the minimum length of a word $a_1 a_2 \cdots a_r \in A^*$ with the property that $g a_1 a_2 \cdots a_r = h$.

The Cayley graph $\Gamma(G, A)$



Vertices: G

Edges: $g \sim h \Leftrightarrow h = ga$ for some $a \in A$

The (?) Cayley graph of a group



The (?) Cayley graph of a group



Conclusion

Changing the finite generating set can result in spaces that are not isometric.

Idea: These two spaces look the same **when viewed from far enough away**. This idea is formalised via the notion of **quasi-isometry**.

Quasi-isometry for metric spaces

Definition

Let (X, d_X) and (Y, d_Y) be two metric spaces. A map $f : X \rightarrow Y$ is a **quasi-isometric embedding** if there exist constants $\lambda \geq 1$ and $C \geq 0$ such that

$$\frac{1}{\lambda}d_X(a, b) - C \leq d_Y(f(a), f(b)) \leq \lambda d_X(a, b) + C,$$

for all $a, b \in X$.

Quasi-isometry for metric spaces

Definition

Let (X, d_X) and (Y, d_Y) be two metric spaces. A map $f : X \rightarrow Y$ is a **quasi-isometric embedding** if there exist constants $\lambda \geq 1$ and $C \geq 0$ such that

$$\frac{1}{\lambda}d_X(a, b) - C \leq d_Y(f(a), f(b)) \leq \lambda d_X(a, b) + C,$$

for all $a, b \in X$. The metric spaces (X, d_X) and (Y, d_Y) are **quasi-isometric** if in addition there is a constant $D \geq 0$ such that every point in Y has a distance at most D from some point in the image $f(X)$.

Quasi-isometry for metric spaces


Definition

Let (X, d_X) and (Y, d_Y) be two metric spaces. A map $f : X \rightarrow Y$ is a **quasi-isometric embedding** if there exist constants $\lambda \geq 1$ and $C \geq 0$ such that

$$\frac{1}{\lambda}d_X(a, b) - C \leq d_Y(f(a), f(b)) \leq \lambda d_X(a, b) + C,$$

for all $a, b \in X$. The metric spaces (X, d_X) and (Y, d_Y) are **quasi-isometric** if in addition there is a constant $D \geq 0$ such that every point in Y has a distance at most D from some point in the image $f(X)$.

- ▶ Quasi-isometry is an **equivalence relation between metric spaces**, which ignores finite details.



Quasi-isometry for finitely generated groups

Proposition

Let A and B be two finite generating sets for the group G .
Then (G, d_A) and (G, d_B) are quasi-isometric.

The quasi-isometry class of a group

Given a finitely generated group G , the metric space (G, d_A) is well defined up to quasi-isometry by the group G alone.

In particular, given two finitely generated groups G and H , one may ask whether they are quasi-isometric or not, without reference to any specific choice of finite generating sets.

Tigers, Lions and Frogs



Tigers and lions look similar and, genetically, they have a lot in common.

Tigers and frogs on the other hand...

Tigers, Lions and Frogs



Tigers and lions look similar and, genetically, they have a lot in common.

Tigers and frogs on the other hand...

Quasi-isometric groups look similar and, algebraically, they have a lot in common.

Quasi-isometry invariants

Quasi-isometry invariants

A property (P) of finitely generated groups is said to be a **quasi-isometry invariant** if, whenever G_1 and G_2 are quasi-isometric finitely generated groups,

$$G_1 \text{ has property } (P) \Leftrightarrow G_2 \text{ has property } (P).$$

Quasi-isometry invariants of groups include being...

(i) Finite; (ii) Infinite virtually cyclic; (iii) Finitely presented; (iv) Virtually abelian; (v) Virtually nilpotent; (vi) Virtually free; (vii) Amenable; (viii) Hyperbolic; (ix) Accessible; (x) Type of growth; (xi) Finitely presented with solvable word problem; (xii) Satisfying the homological finiteness condition F_n or the condition FP_n ; (xiii) Number of ends.


General philosophy

Algebra	Combinatorics	Geometry
Groups	Graphs	Metric spaces
Monoids / Semigroups	Digraphs	??

?? = directed metric spaces = semimetric spaces

Semimetric space = a set equipped with an asymmetric, partially-defined distance function.

Cayley graphs of semigroups and monoids



The bicyclic monoid $B = \langle b, c \mid bc = 1 \rangle$

Semimetric spaces

Definition (Semimetric space)

A **semimetric space** is a pair (X, d) where X is a set, and

$$d : X \times X \rightarrow \mathbb{R}^\infty = \mathbb{R}^{\geq 0} \cup \{\infty\}$$

is a function satisfying:

(i) $d(x, y) = 0$ if and only if $x = y$; and

(ii) $d(x, z) \leq d(x, y) + d(y, z)$;

for all $x, y, z \in X$.

Here $\mathbb{R}^\infty = \mathbb{R}^{\geq 0} \cup \{\infty\}$ with the obvious order, and we set

$$\infty + x = x + \infty = y\infty = \infty y = \infty$$

for all $x \in \mathbb{R}^\infty$ and $y \in \mathbb{R}^\infty \setminus \{0\}$.

Monoids as semimetric spaces

M - monoid generated by a finite set A .

M gives rise to a semimetric space (M, d_A) with word semimetric d_A .

Points: M

Directed distance: $d_A(x, y)$ the minimum length of a word $a_1 a_2 \cdots a_r \in A^*$ with the property that $xa_1 a_2 \cdots a_r = y$, or ∞ if there is no such word.

The (right) Cayley graph $\Gamma(M, A)$

Vertices: M

Directed edges: $x \rightarrow y \Leftrightarrow y = xa$ for some $a \in A$

Quasi-isometry for semimetric spaces

Definition

Let (X, d_X) and (Y, d_Y) be two semimetric spaces. A map $f : X \rightarrow Y$ is a **quasi-isometric embedding** if there exist constants $\lambda \geq 1$ and $C \geq 0$ such that

$$\frac{1}{\lambda}d_X(a, b) - C \leq d_Y(f(a), f(b)) \leq \lambda d_X(a, b) + C,$$

for all $a, b \in X$.

Quasi-isometry for semimetric spaces

Definition

Let (X, d_X) and (Y, d_Y) be two semimetric spaces. A map $f : X \rightarrow Y$ is a **quasi-isometric embedding** if there exist constants $\lambda \geq 1$ and $C \geq 0$ such that


$$\frac{1}{\lambda}d_X(a, b) - C \leq d_Y(f(a), f(b)) \leq \lambda d_X(a, b) + C,$$

for all $a, b \in X$.

The semimetric spaces (X, d_X) and (Y, d_Y) are **quasi-isometric** if in addition there is a constant $D \geq 0$ such that for every $y \in Y$ there exists a $z \in f(X)$ such that

$$d_Y(y, z) \leq D, \quad \text{and} \quad d_Y(z, y) \leq D.$$

- ▶ Quasi-isometry is an equivalence relation between semimetric spaces.



Quasi-isometry for finitely generated monoids

Proposition

Let A and B be two finite generating sets for a monoid M .

Then the semimetric space (M, d_A) is quasi-isometric to the semimetric space (M, d_B) .

The quasi-isometry class of a monoid

The semimetric space (M, d_A) is well defined up to quasi-isometry by the finitely generated monoid M alone.

In particular, given two finitely generated monoids M and N , one may ask whether they are quasi-isometric or not, without reference to any specific choice of finite generating sets.

Quasi-isometry invariants of monoids

Theorem

The following properties are all quasi-isometry invariants of finitely generated monoids:

- ▶ Finiteness, Number of right ideals;
- ▶ Being a group (for monoids), being right simple (for semigroups);
- ▶ Type of growth;
- ▶ Number of ends (in the sense of [Jackson and Kilibarda \(2009\)](#)).

However...

There are a number of important properties which are quasi-isometry invariants of groups, for which it is currently not known whether they are quasi-isometry invariants of monoids.

Finite presentability and the word problem

Two open problems

1. Is **finite presentability** a quasi-isometry invariant of finitely generated monoids?
2. Is being **finitely presented with solvable word problem** a quasi-isometry invariant of finitely generated monoids?

Idea: Can anything be said for classes of monoids that lie between monoids and groups?

Consider monoids that are:

1. Left cancellative;
2. Have finitely many left and right ideals.

Obviously any group satisfies both (1) and (2).

Left cancellative semigroups and monoids

Left cancellativity: $ab = ac \Rightarrow b = c$.

Right cancellativity, and cancellativity are defined analogously.

Interesting classes of cancellative monoids

- ▶ Divisibility monoids (Droste & Kuske (2001));
- ▶ Garside monoids; includes, spherical Artin monoids, Braid monoids of complex reflection groups etc. (Dehornoy & Paris (1999)).

One-relator monoids

- ▶ Adyan and Oganesyan (1987): Decidability of the word problem for one relator monoids is reducible to the left cancellative case.
- ▶ Motivates the development of new methods for approaching the word problem for finitely presented left cancellative monoids.

Directed 2-complexes

Directed graph

A **digraph** Γ consists of:

V - vertices, E - directed edges, and functions

$\iota, \tau: E \rightarrow V$, expressing the initial / terminal vertices of each directed edge.

A path in Γ is a sequence of composable directed edges $p = e_1 e_2 \dots e_r$

ι and τ extend to paths in the obvious way.

$P = P(\Gamma)$ - set of all paths from Γ

$p, q \in P$ are parallel, written $p \parallel q$, if $\iota p = \iota q$ and $\tau p = \tau q$.

Directed 2-complexes

Directed graph

A **digraph** Γ consists of:

V - vertices, E - directed edges, and functions

$\iota, \tau: E \rightarrow V$, expressing the initial / terminal vertices of each directed edge.

A path in Γ is a sequence of composable directed edges $p = e_1 e_2 \dots e_r$
 ι and τ extend to paths in the obvious way.

$P = P(\Gamma)$ - set of all paths from Γ

$p, q \in P$ are parallel, written $p \parallel q$, if $\iota p = \iota q$ and $\tau p = \tau q$.


Directed 2-complex (following Guba & Sapir (2006))

Γ - digraph, together with F - set of **2-cells**, and maps

$[\cdot]: F \rightarrow P$, $[\cdot]: F \rightarrow P$, and $^{-1}: F \rightarrow F$ called **top**, **bottom**, and **inverse** such that

- ▶ for every $f \in F$, the paths $[f]$ and $[f]$ are parallel;
- ▶ $^{-1}$ is an involution without fixed points, and $[f^{-1}] = [f]$, $[f^{-1}] = [f]$ for every $f \in F$.

Directed 2-complex




2-paths in directed 2-complexes

K - a directed 2-complex, with underlying digraph Γ , and set of faces F
The 1-paths in K are the paths in Γ .

Definition (2-path)

An **atomic** 2-path δ is a triple (p, f, q) where p, q are 1-paths, $f \in F$ and:



Define $[\delta] = p [f] q$ and $[\delta] = p [f] q$.

A **2-path** in K is then a sequence $\delta = \delta_1 \delta_2 \dots \delta_n$ of composable atomic 2-paths, meaning $[\delta_i] = [\delta_{i+1}]$ for all i .

Define $[\delta] = [\delta_1]$ and $[\delta] = [\delta_n]$ - the top and the bottom of the 2-path δ .

Directed homotopy and simple connectedness

Directed homotopy

K - directed 2-complex, 1-paths p, q in K are **homotopic** if there is a 2-path δ in K such that $[\delta] = p$ and $[\delta] = q$.

K is **directed simply connected** if for every pair $p \parallel q$ of parallel paths, p and q are homotopic in K .

Directed homotopy and simple connectedness

Directed homotopy

K - directed 2-complex, 1-paths p, q in K are **homotopic** if there is a 2-path δ in K such that $[\delta] = p$ and $[\delta] = q$.

K is **directed simply connected** if for every pair $p \parallel q$ of parallel paths, p and q are homotopic in K .

Quasi-simple connectedness

Γ - digraph, $n \in \mathbb{N}$

$K_n(\Gamma)$ = directed 2-complex with underlying digraph Γ and face set


$$F = \{(p, q) \mid p \text{ and } q \text{ are parallel paths in } \Gamma \text{ with } |p| + |q| \leq n\}$$

and $[(p, q)] = p$, $[(p, q)] = q$ and $(p, q)^{-1} = (q, p)$.

Note: $K_n(\Gamma)$ is the natural **directed analogue of the Rips complex**.

- ▶ We say Γ is **quasi-simply-connected** if $K_n(\Gamma)$ is directed simply connected for some n .

Directed 2-complexes




Consider the 2-complex $K_4(\Gamma)$ where
 Γ - right Cayley graph of the monoid $\langle a, b \mid ab = ba \rangle$.
 Diagram illustrates a 2-path δ of length 4 in $K_4(\Gamma)$ with

$$[\delta] = aaabba, \quad \text{and} \quad [\delta] = abbaaa.$$

Observe: This 2-path corresponds to a derivation of the equivalence $aaabba = abbaaa$ in the monoid.

Directed 2-complexes




Consider the 2-complex $K_4(\Gamma)$ where
 Γ - right Cayley graph of the monoid $\langle a, b \mid ab = ba \rangle$.
Diagram illustrates a 2-path δ of length 4 in $K_4(\Gamma)$ with

$$[\delta] = aaabba, \quad \text{and} \quad [\delta] = abbaaa.$$

Observe: This 2-path corresponds to a derivation of the equivalence $aaabba = abbaaa$ in the monoid.

Directed 2-complexes




Consider the 2-complex $K_4(\Gamma)$ where
 Γ - right Cayley graph of the monoid $\langle a, b \mid ab = ba \rangle$.
Diagram illustrates a 2-path δ of length 4 in $K_4(\Gamma)$ with

$$[\delta] = aaabba, \quad \text{and} \quad [\delta] = abbaaa.$$

Observe: This 2-path corresponds to a derivation of the equivalence $aaabba = abbaaa$ in the monoid.

Directed 2-complexes




Consider the 2-complex $K_4(\Gamma)$ where
 Γ - right Cayley graph of the monoid $\langle a, b \mid ab = ba \rangle$.
Diagram illustrates a 2-path δ of length 4 in $K_4(\Gamma)$ with

$$[\delta] = aaabba, \quad \text{and} \quad [\delta] = abbaaa.$$

Observe: This 2-path corresponds to a derivation of the equivalence $aaabba = abbaaa$ in the monoid.

Directed 2-complexes




Consider the 2-complex $K_4(\Gamma)$ where
 Γ - right Cayley graph of the monoid $\langle a, b \mid ab = ba \rangle$.
 Diagram illustrates a 2-path δ of length 4 in $K_4(\Gamma)$ with

$$[\delta] = aaabba, \quad \text{and} \quad [\delta] = abbaaa.$$

Observe: This 2-path corresponds to a derivation of the equivalence $aaabba = abbaaa$ in the monoid.

Directed 2-complexes



Consider the 2-complex $K_4(\Gamma)$ where
 Γ - right Cayley graph of the monoid $\langle a, b \mid ab = ba \rangle$.
Diagram illustrates a 2-path δ of length 4 in $K_4(\Gamma)$ with

$$[\delta] = aaabba, \quad \text{and} \quad [\delta] = abbaaa.$$

Observe: This 2-path corresponds to a derivation of the equivalence $aaabba = abbaaa$ in the monoid.

Finite presentability and the word problem

Theorem

Let S be a left cancellative monoid generated by a finite set A . Then:

- S is finitely presented $\Leftrightarrow \Gamma(S, A)$ is quasi-simply-connected.

Proposition

The property of being quasi-simply-connected is a quasi-isometry invariant of directed graphs.

Theorem

Let M and N be left cancellative, finitely generated monoids which are quasi-isometric. Then M is finitely presentable $\Leftrightarrow N$ is finitely presentable.

Finite presentability and the word problem

Theorem

Let S be a left cancellative monoid generated by a finite set A . Then:

- S is finitely presented $\Leftrightarrow \Gamma(S, A)$ is quasi-simply-connected.

Proposition

The property of being quasi-simply-connected is a quasi-isometry invariant of directed graphs.

Theorem

Let M and N be left cancellative, finitely generated monoids which are quasi-isometric. Then M is finitely presentable $\Leftrightarrow N$ is finitely presentable.

By defining and studying [Dehn functions of directed 2-complexes](#) and their behaviour under quasi-isometry, one can show:

Theorem

Let M and N be left cancellative, finitely presentable monoids which are quasi-isometric. Then M has solvable word problem if and only if N has solvable word problem.

Monoids with finitely many left and right ideals


We established a **Švarc–Milnor Lemma for groups acting on geodesic semimetric spaces**. Applying this result to Schützenberger groups acting on Schützenberger graphs leads to the following.

Theorem

Let M be a finitely generated monoid with finitely many left and right ideals. Then M is finitely presented if and only if all right Schützenberger graphs of M are quasi-simply-connected.

Theorem

For finitely generated monoids with finitely many left and right ideals, finite presentability is a quasi-isometry invariant.



- Analogous result for finitely presented with solvable word problem holds.

Monoids in general

Conclusion

For certain spacial classes of monoids quasi-simple-connectedness of directed 2-complexes can be used to capture geometrically the property of being finitely presented.

For finitely generated monoids in general, quasi-simple-connectedness is far from capturing finite presentability.

Indeed, in general:

1. Finite presentability \Rightarrow Quasi-simple-connectedness, and
2. Quasi-simple-connectedness \Rightarrow Finite presentability.

Monoids in general

Quasi-simple-connectedness \Rightarrow Finite presentability

Example


$$\mathbb{N} = \{0, 1, 2, \dots\}, \quad \emptyset \subsetneq X \subsetneq \mathbb{N}$$

$$M(X) = \langle a, b, c, d, e \mid ab^i c = ab^i d \ (i \in X), \quad ab^j c = ab^j e \ (j \notin X) \rangle.$$

- (i) $M(X)$ does not admit a finite presentation.
- (ii) The word problem for $M(X)$ is solvable $\Leftrightarrow X$ is a recursive subset of \mathbb{N} .
- (iii) For any subsets X and Y of \mathbb{N} , the semigroups $M(X)$ and $M(Y)$ are isometric to each other, and to a directed rooted tree in which every vertex has out-degree 4 or 5.
- (iv) The Cayley graph of $M(X)$ is quasi-simply connected, since it is a tree.

Consequences

1. Quasi-simple-connectedness \Rightarrow Finite presentability.
2. Having solvable word problem is **not** a quasi-isometry invariant of finitely generated monoids.



Future directions

- ▶ For arbitrary finitely generated monoids decide whether the properties of being
 - (a) finitely presented;
 - (b) finitely presented with solvable word problem,are quasi-isometry invariants.
- ▶ Are they isometry invariants?
- ▶ Are there other natural classes of monoids for which (a) and (b) are quasi-isometry invariants?
- ▶ Investigate other properties from the point of view of quasi-isometry (e.g. Amenable semigroups (Day (1957)) / Følner conditions in digraphs).
- ▶ We have restricted our attention to the geometry of **right Cayley graphs** only. What if one considers the geometry of right *and* left Cayley graphs?