

Islands and proximity domains

Eszter K. Horváth, Szeged

Co-authors: Stephan Foldes, Sándor Radeleczki, Tamás Waldhauser

Novi Sad, 2013, June 5.

$$U \in \mathcal{C} \subseteq \mathcal{K} \subseteq \mathcal{P}(U)$$

Let $h: U \rightarrow \mathbb{R}$ be a height function and let $S \in \mathcal{C}$ be a nonempty set.

We denote the cover relation of the poset (\mathcal{K}, \subseteq) by \prec , and we write $K_1 \preceq K_2$ if $K_1 \prec K_2$ or $K_1 = K_2$.

We say that S is a *island* with respect to the triple $(\mathcal{C}, \mathcal{K}, h)$, if every $K \in \mathcal{K}$ with $S \prec K$ satisfies

$$h(u) < \min h(S) \text{ for all } u \in K \setminus S.$$

Island domain

$$U \in \mathcal{C} \subseteq \mathcal{K} \subseteq \mathcal{P}(U)$$

Let $h: U \rightarrow \mathbb{R}$ be a height function and let $S \in \mathcal{C}$ be a nonempty set.

We denote the cover relation of the poset (\mathcal{K}, \subseteq) by \prec , and we write $K_1 \preceq K_2$ if $K_1 \prec K_2$ or $K_1 = K_2$.

We say that S is a *island* with respect to the triple $(\mathcal{C}, \mathcal{K}, h)$, if every $K \in \mathcal{K}$ with $S \prec K$ satisfies

$$h(u) < \min h(S) \text{ for all } u \in K \setminus S.$$

$$U \in \mathcal{C} \subseteq \mathcal{K} \subseteq \mathcal{P}(U)$$

Let $h: U \rightarrow \mathbb{R}$ be a height function and let $S \in \mathcal{C}$ be a nonempty set.

We denote the cover relation of the poset (\mathcal{K}, \subseteq) by \prec , and we write $K_1 \preceq K_2$ if $K_1 \prec K_2$ or $K_1 = K_2$.

We say that S is a *island* with respect to the triple $(\mathcal{C}, \mathcal{K}, h)$, if every $K \in \mathcal{K}$ with $S \prec K$ satisfies

$$h(u) < \min h(S) \text{ for all } u \in K \setminus S.$$

„Closeness” relation

$(\mathcal{C}, \mathcal{K})$

$\delta \subseteq \mathcal{C} \times \mathcal{C}$

$$A\delta B \Leftrightarrow \exists K \in \mathcal{K} : A \preceq K \text{ and } K \cap B \neq \emptyset. \quad (1)$$

It is easy to verify that relation δ satisfies the following properties for all $A, B, C \in \mathcal{C}$ whenever $B \cup C \in \mathcal{C}$:

$$A\delta B \Rightarrow B \neq \emptyset;$$

$$A \cap B \neq \emptyset \Rightarrow A\delta B;$$

$$A\delta(B \cup C) \Leftrightarrow (A\delta B \text{ or } A\delta C).$$

„Closeness” relation

$(\mathcal{C}, \mathcal{K})$

$\delta \subseteq \mathcal{C} \times \mathcal{C}$

$$A\delta B \Leftrightarrow \exists K \in \mathcal{K} : A \preceq K \text{ and } K \cap B \neq \emptyset. \quad (1)$$

It is easy to verify that relation δ satisfies the following properties for all $A, B, C \in \mathcal{C}$ whenever $B \cup C \in \mathcal{C}$:

$$A\delta B \Rightarrow B \neq \emptyset;$$

$$A \cap B \neq \emptyset \Rightarrow A\delta B;$$

$$A\delta(B \cup C) \Leftrightarrow (A\delta B \text{ or } A\delta C).$$

„Closeness” relation

$(\mathcal{C}, \mathcal{K})$

$\delta \subseteq \mathcal{C} \times \mathcal{C}$

$$A\delta B \Leftrightarrow \exists K \in \mathcal{K}: A \preceq K \text{ and } K \cap B \neq \emptyset. \quad (1)$$

It is easy to verify that relation δ satisfies the following properties for all $A, B, C \in \mathcal{C}$ whenever $B \cup C \in \mathcal{C}$:

$$A\delta B \Rightarrow B \neq \emptyset;$$

$$A \cap B \neq \emptyset \Rightarrow A\delta B;$$

$$A\delta(B \cup C) \Leftrightarrow (A\delta B \text{ or } A\delta C).$$

„Closeness” relation

$(\mathcal{C}, \mathcal{K})$

$\delta \subseteq \mathcal{C} \times \mathcal{C}$

$$A\delta B \Leftrightarrow \exists K \in \mathcal{K} : A \preceq K \text{ and } K \cap B \neq \emptyset. \quad (1)$$

It is easy to verify that relation δ satisfies the following properties for all $A, B, C \in \mathcal{C}$ whenever $B \cup C \in \mathcal{C}$:

$$A\delta B \Rightarrow B \neq \emptyset;$$

$$A \cap B \neq \emptyset \Rightarrow A\delta B;$$

$$A\delta(B \cup C) \Leftrightarrow (A\delta B \text{ or } A\delta C).$$

We say that $A, B \in \mathcal{C}$ are *distant* if neither $A\delta B$ nor $B\delta A$ holds.

It is easy to see that in this case A and B are also incomparable (in fact, disjoint), whenever $A, B \neq \emptyset$.

A nonempty family $\mathcal{H} \subseteq \mathcal{C}$ will be called a *distant family*, if any two incomparable members of \mathcal{H} are distant.

Lemma If $\mathcal{H} \subseteq \mathcal{C}$ is a distant family, then \mathcal{H} is CDW-independent. Moreover, if $U \in \mathcal{H}$, then U is admissible.

We say that $A, B \in \mathcal{C}$ are *distant* if neither $A\delta B$ nor $B\delta A$ holds.

It is easy to see that in this case A and B are also incomparable (in fact, disjoint), whenever $A, B \neq \emptyset$.

A nonempty family $\mathcal{H} \subseteq \mathcal{C}$ will be called a *distant family*, if any two incomparable members of \mathcal{H} are distant.

Lemma If $\mathcal{H} \subseteq \mathcal{C}$ is a distant family, then \mathcal{H} is CDW-independent. Moreover, if $U \in \mathcal{H}$, then U is admissible.

We say that $A, B \in \mathcal{C}$ are *distant* if neither $A\delta B$ nor $B\delta A$ holds.

It is easy to see that in this case A and B are also incomparable (in fact, disjoint), whenever $A, B \neq \emptyset$.

A nonempty family $\mathcal{H} \subseteq \mathcal{C}$ will be called a *distant family*, if any two incomparable members of \mathcal{H} are distant.

Lemma If $\mathcal{H} \subseteq \mathcal{C}$ is a distant family, then \mathcal{H} is CDW-independent. Moreover, if $U \in \mathcal{H}$, then U is admissible.

We say that $A, B \in \mathcal{C}$ are *distant* if neither $A\delta B$ nor $B\delta A$ holds.

It is easy to see that in this case A and B are also incomparable (in fact, disjoint), whenever $A, B \neq \emptyset$.

A nonempty family $\mathcal{H} \subseteq \mathcal{C}$ will be called a *distant family*, if any two incomparable members of \mathcal{H} are distant.

Lemma If $\mathcal{H} \subseteq \mathcal{C}$ is a distant family, then \mathcal{H} is CDW-independent. Moreover, if $U \in \mathcal{H}$, then U is admissible.

Definitions

Let $\mathbb{P} = (P, \leq)$ be a partially ordered set and $a, b \in P$. The elements a and b are called *disjoint* and we write $a \perp b$ if

either \mathbb{P} has least element $0 \in P$ and $\inf\{a, b\} = 0$,

or \mathbb{P} is without 0 , then a and b have no common lower bound.

A nonempty set $X \subseteq P$ is called *CD-independent* if for any $x, y \in X$, $x \leq y$ or $y \leq x$ or $x \perp y$ holds.

Definitions

Let $\mathbb{P} = (P, \leq)$ be a partially ordered set and $a, b \in P$. The elements a and b are called *disjoint* and we write $a \perp b$ if

either \mathbb{P} has least element $0 \in P$ and $\inf\{a, b\} = 0$,

or if \mathbb{P} is without 0 , then a and b have no common lowerbound.

A nonempty set $X \subseteq P$ is called *CD-independent* if for any $x, y \in X$, $x \leq y$ or $y \leq x$ or $x \perp y$ holds.

Definitions

Let $\mathbb{P} = (P, \leq)$ be a partially ordered set and $a, b \in P$. The elements a and b are called *disjoint* and we write $a \perp b$ if

either \mathbb{P} has least element $0 \in P$ and $\inf\{a, b\} = 0$,

or if \mathbb{P} is without 0 , then a and b have no common lowerbound.

A nonempty set $X \subseteq P$ is called *CD-independent* if for any $x, y \in X$, $x \leq y$ or $y \leq x$ or $x \perp y$ holds.

Definitions

Let $\mathbb{P} = (P, \leq)$ be a partially ordered set and $a, b \in P$. The elements a and b are called *disjoint* and we write $a \perp b$ if

either \mathbb{P} has least element $0 \in P$ and $\inf\{a, b\} = 0$,

or if \mathbb{P} is without 0 , then a and b have no common lowerbound.

A nonempty set $X \subseteq P$ is called *CD-independent* if for any $x, y \in X$, $x \leq y$ or $y \leq x$ or $x \perp y$ holds.

Definitions

Let $\mathbb{P} = (P, \leq)$ be a partially ordered set and $a, b \in P$. The elements a and b are called *disjoint* and we write $a \perp b$ if

either \mathbb{P} has least element $0 \in P$ and $\inf\{a, b\} = 0$,

or if \mathbb{P} is without 0 , then a and b have no common lowerbound.

A nonempty set $X \subseteq P$ is called *CD-independent* if for any $x, y \in X$, $x \leq y$ or $y \leq x$ or $x \perp y$ holds.

Definition A family $\mathcal{H} \subseteq \mathcal{P}(U)$ is *weakly independent* if

$$H \subseteq \bigcup_{i \in I} H_i \implies \exists i \in I : H \subseteq H_i \quad (2)$$

holds for all $H \in \mathcal{H}, H_i \in \mathcal{H} (i \in I)$. If \mathcal{H} is both CD-independent and weakly independent, then we say that \mathcal{H} is *CDW-independent*.

Definition

Let $\mathcal{H} \subseteq \mathcal{C} \setminus \{\emptyset\}$ be a family of sets such that $U \in \mathcal{H}$. We say that \mathcal{H} is *admissible*, if for every nonempty antichain $\mathcal{A} \subseteq \mathcal{H}$


$$\exists H \in \mathcal{A} \forall K \in \mathcal{K} : H \subset K \implies K \notin \bigcup \mathcal{A}. \quad (3)$$

Connective island domains

Definition

A pair $(\mathcal{C}, \mathcal{K})$ is a *connective island domain* if

$$\forall A, B \in \mathcal{C} : (A \cap B \neq \emptyset \text{ and } B \not\subseteq A) \implies \exists K \in \mathcal{K} : A \subset K \subseteq A \cup B.$$



Theorem

The following three conditions are equivalent for any pair $(\mathcal{C}, \mathcal{K})$:

(i) $(\mathcal{C}, \mathcal{K})$ is a connective island domain.

(ii) Every system of pre-islands corresponding to $(\mathcal{C}, \mathcal{K})$ is CD-independent.

(iii) Every system of pre-islands corresponding to $(\mathcal{C}, \mathcal{K})$ is CDW-independent.

Theorem

The following three conditions are equivalent for any pair $(\mathcal{C}, \mathcal{K})$:

(i) $(\mathcal{C}, \mathcal{K})$ is a connective island domain.

(ii) Every system of pre-islands corresponding to $(\mathcal{C}, \mathcal{K})$ is CD-independent.

(iii) Every system of pre-islands corresponding to $(\mathcal{C}, \mathcal{K})$ is CDW-independent.

Theorem

The following three conditions are equivalent for any pair $(\mathcal{C}, \mathcal{K})$:

(i) $(\mathcal{C}, \mathcal{K})$ is a connective island domain.

(ii) Every system of pre-islands corresponding to $(\mathcal{C}, \mathcal{K})$ is CD-independent.


(iii) Every system of pre-islands corresponding to $(\mathcal{C}, \mathcal{K})$ is CDW-independent.

Standard height function

Let us consider a CD-independent family \mathcal{H} .

Clearly, for every $u \in U$, the set of members of \mathcal{H} containing u is a finite chain.

The *standard height function* of \mathcal{H} assigns to each element u the length of this chain, i.e., one less than the number of members of \mathcal{H} that contain u .




Standard height function

Let us consider a CD-independent family \mathcal{H} .

Clearly, for every $u \in U$, the set of members of \mathcal{H} containing u is a finite chain.

The *standard height function* of \mathcal{H} assigns to each element u the length of this chain, i.e., one less than the number of members of \mathcal{H} that contain u .




Standard height function

Let us consider a CD-independent family \mathcal{H} .

Clearly, for every $u \in U$, the set of members of \mathcal{H} containing u is a finite chain.

The *standard height function* of \mathcal{H} assigns to each element u the length of this chain, i.e., one less than the number of members of \mathcal{H} that contain u .




Standard height function

Let us consider a CD-independent family \mathcal{H} .

Clearly, for every $u \in U$, the set of members of \mathcal{H} containing u is a finite chain.

The *standard height function* of \mathcal{H} assigns to each element u the length of this chain, i.e., one less than the number of members of \mathcal{H} that contain u .



Theorem

Let $(\mathcal{C}, \mathcal{K})$ be a connective island domain and let $\mathcal{H} \subseteq \mathcal{C} \setminus \{\emptyset\}$ with $U \in \mathcal{H}$. If \mathcal{H} is a distant family, then \mathcal{H} is a system of islands; moreover, \mathcal{H} is the system of islands corresponding to its standard height function.

Islands and proximity domains

The island domain $(\mathcal{C}, \mathcal{K})$ is called a *proximity domain*, if it is a connective island domain and the relation δ is symmetric for nonempty sets, that is

$$\forall A, B \in \mathcal{C} \setminus \{\emptyset\} : A\delta B \Leftrightarrow B\delta A. \quad (4)$$

If a relation δ defined on $\mathcal{P}(U)$ satisfies the mentioned three properties and δ is symmetric for nonempty sets, then (U, δ) is called a *proximity space*.

δ satisfies the following properties for all $A, B, C \in \mathcal{C}$ whenever $B \cup C \in \mathcal{C}$:

$$\begin{aligned} A\delta B &\Rightarrow B \neq \emptyset; \\ A \cap B \neq \emptyset &\Rightarrow A\delta B; \\ A\delta(B \cup C) &\Leftrightarrow (A\delta B \text{ or } A\delta C). \end{aligned}$$

The notion goes back to Frigyes Riesz (1908), however this axiomatization is due to Vadim A. Efremovich.

Islands and proximity domains

The island domain $(\mathcal{C}, \mathcal{K})$ is called a *proximity domain*, if it is a connective island domain and the relation δ is symmetric for nonempty sets, that is

$$\forall A, B \in \mathcal{C} \setminus \{\emptyset\} : A\delta B \Leftrightarrow B\delta A. \quad (4)$$

If a relation δ defined on $\mathcal{P}(U)$ satisfies the mentioned three properties and δ is symmetric for nonempty sets, then (U, δ) is called a *proximity space*.

δ satisfies the following properties for all $A, B, C \in \mathcal{C}$ whenever $B \cup C \in \mathcal{C}$:

$$\begin{aligned} A\delta B &\Rightarrow B \neq \emptyset; \\ A \cap B \neq \emptyset &\Rightarrow A\delta B; \\ A\delta(B \cup C) &\Leftrightarrow (A\delta B \text{ or } A\delta C). \end{aligned}$$

The notion goes back to Frigyes Riesz (1908), however this axiomatization is due to Vadim A. Efremovich.

Islands and proximity domains

The island domain $(\mathcal{C}, \mathcal{K})$ is called a *proximity domain*, if it is a connective island domain and the relation δ is symmetric for nonempty sets, that is

$$\forall A, B \in \mathcal{C} \setminus \{\emptyset\} : A\delta B \Leftrightarrow B\delta A. \quad (4)$$

If a relation δ defined on $\mathcal{P}(U)$ satisfies the mentioned three properties and δ is symmetric for nonempty sets, then (U, δ) is called a *proximity space*.

δ satisfies the following properties for all $A, B, C \in \mathcal{C}$ whenever $B \cup C \in \mathcal{C}$:

$$\begin{aligned} A\delta B &\Rightarrow B \neq \emptyset; \\ A \cap B \neq \emptyset &\Rightarrow A\delta B; \\ A\delta(B \cup C) &\Leftrightarrow (A\delta B \text{ or } A\delta C). \end{aligned}$$

The notion goes back to Frigyes Riesz (1908), however this axiomatization is due to Vadim A. Efremovich.

Islands and proximity domains

The island domain $(\mathcal{C}, \mathcal{K})$ is called a *proximity domain*, if it is a connective island domain and the relation δ is symmetric for nonempty sets, that is

$$\forall A, B \in \mathcal{C} \setminus \{\emptyset\} : A\delta B \Leftrightarrow B\delta A. \quad (4)$$

If a relation δ defined on $\mathcal{P}(U)$ satisfies the mentioned three properties and δ is symmetric for nonempty sets, then (U, δ) is called a *proximity space*.

δ satisfies the following properties for all $A, B, C \in \mathcal{C}$ whenever $B \cup C \in \mathcal{C}$:

$$\begin{aligned} A\delta B &\Rightarrow B \neq \emptyset; \\ A \cap B \neq \emptyset &\Rightarrow A\delta B; \\ A\delta(B \cup C) &\Leftrightarrow (A\delta B \text{ or } A\delta C). \end{aligned}$$

The notion goes back to Frigyes Riesz (1908), however this axiomatization is due to Vadim A. Efremovich.

Proposition

If $(\mathcal{C}, \mathcal{K})$ is a proximity domain, then any system of islands corresponding to $(\mathcal{C}, \mathcal{K})$ is a distant system.

Proof

$$h(b) < \min h(A) \leq h(a)$$

$$h(a) < \min h(B) \leq h(b)$$

Proposition

If $(\mathcal{C}, \mathcal{K})$ is a proximity domain, then any system of islands corresponding to $(\mathcal{C}, \mathcal{K})$ is a distant system.

Proof

$$h(b) < \min h(A) \leq h(a)$$

$$h(a) < \min h(B) \leq h(b)$$

Characterization for system of islands for proximity domains

Corollary

If $(\mathcal{C}, \mathcal{K})$ is a proximity domain, and $\mathcal{H} \subseteq \mathcal{C} \setminus \{\emptyset\}$ with $U \in \mathcal{H}$, then \mathcal{H} is a system of islands if and only if \mathcal{H} is a distant family. Moreover, in this case \mathcal{H} is the system of islands corresponding to its standard height function.

$$U \in \mathcal{C} \subseteq \mathcal{K} \subseteq \mathcal{P}(U)$$

Let $h: U \rightarrow \mathbb{R}$ be a height function and let $S \in \mathcal{C}$ be a nonempty set.

We say that S is an *pre-island* with respect to the triple $(\mathcal{C}, \mathcal{K}, h)$, if every $K \in \mathcal{K}$ with $S \prec K$ satisfies

$$\min h(K) < \min h(S).$$

We say that S is a *island* with respect to the triple $(\mathcal{C}, \mathcal{K}, h)$, if every $K \in \mathcal{K}$ with $S \prec K$ satisfies

$$h(u) < \min h(S) \text{ for all } u \in K \setminus S.$$

$$U \in \mathcal{C} \subseteq \mathcal{K} \subseteq \mathcal{P}(U)$$

Let $h: U \rightarrow \mathbb{R}$ be a height function and let $S \in \mathcal{C}$ be a nonempty set.

We say that S is an *pre-island* with respect to the triple $(\mathcal{C}, \mathcal{K}, h)$, if every $K \in \mathcal{K}$ with $S \prec K$ satisfies

$$\min h(K) < \min h(S).$$

We say that S is a *island* with respect to the triple $(\mathcal{C}, \mathcal{K}, h)$, if every $K \in \mathcal{K}$ with $S \prec K$ satisfies

$$h(u) < \min h(S) \text{ for all } u \in K \setminus S.$$

Example

Let A_1, \dots, A_n be nonempty sets, and let $\mathcal{I} \subseteq A_1 \times \dots \times A_n$. Let us define

$$U = A_1 \times \dots \times A_n,$$

$$\mathcal{K} = \{B_1 \times \dots \times B_n : \emptyset \neq B_i \subseteq A_i, 1 \leq i \leq n\}$$

$$\mathcal{C} = \{C \in \mathcal{K} : C \subseteq \mathcal{I}\} \cup \{U\},$$

and let $h: U \rightarrow \{0, 1\}$ be the height function given by

$$h(a_1, \dots, a_n) := \begin{cases} 1, & \text{if } (a_1, \dots, a_n) \in \mathcal{I}; \\ 0, & \text{if } (a_1, \dots, a_n) \in U \setminus \mathcal{I}; \end{cases} \quad \text{for all } (a_1, \dots, a_n) \in U.$$

It is easy to see that the pre-islands corresponding to the triple $(\mathcal{C}, \mathcal{K}, h)$ are exactly U and the maximal elements of the poset $(\mathcal{C} \setminus \{U\}, \subseteq)$.

formal concepts

prime implicants of a Boolean function

Example

Let A_1, \dots, A_n be nonempty sets, and let $\mathcal{I} \subseteq A_1 \times \dots \times A_n$. Let us define

$$U = A_1 \times \dots \times A_n,$$

$$\mathcal{K} = \{B_1 \times \dots \times B_n : \emptyset \neq B_i \subseteq A_i, 1 \leq i \leq n\}$$

$$\mathcal{C} = \{C \in \mathcal{K} : C \subseteq \mathcal{I}\} \cup \{U\},$$

and let $h: U \rightarrow \{0, 1\}$ be the height function given by

$$h(a_1, \dots, a_n) := \begin{cases} 1, & \text{if } (a_1, \dots, a_n) \in \mathcal{I}; \\ 0, & \text{if } (a_1, \dots, a_n) \in U \setminus \mathcal{I}; \end{cases} \quad \text{for all } (a_1, \dots, a_n) \in U.$$

It is easy to see that the pre-islands corresponding to the triple $(\mathcal{C}, \mathcal{K}, h)$ are exactly U and the maximal elements of the poset $(\mathcal{C} \setminus \{U\}, \subseteq)$.

formal concepts

prime implicants of a Boolean function

Definition

Let $\mathcal{H} \subseteq \mathcal{C} \setminus \{\emptyset\}$ be a family of sets such that $U \in \mathcal{H}$. We say that \mathcal{H} is *admissible*, if for every nonempty antichain $\mathcal{A} \subseteq \mathcal{H}$

$$\exists H \in \mathcal{A} \forall K \in \mathcal{K} : H \subset K \implies K \notin \bigcup \mathcal{A}. \quad (5)$$

Proposition

Every system of pre-islands is admissible.

Definition

Let $\mathcal{H} \subseteq \mathcal{C} \setminus \{\emptyset\}$ be a family of sets such that $U \in \mathcal{H}$. We say that \mathcal{H} is *admissible*, if for every nonempty antichain $\mathcal{A} \subseteq \mathcal{H}$

$$\exists H \in \mathcal{A} \forall K \in \mathcal{K} : H \subset K \implies K \notin \bigcup \mathcal{A}. \quad (5)$$

Proposition

Every system of pre-islands is admissible.

Theorem

A subfamily of \mathcal{C} is a maximal system of pre-islands if and only if it is a maximal admissible family.

Finally, let us consider the following condition on $(\mathcal{C}, \mathcal{K})$, which is stronger than that of being a connective island domain:

$$\forall K_1, K_2 \in \mathcal{K} : K_1 \cap K_2 \neq \emptyset \implies K_1 \cup K_2 \in \mathcal{K}. \quad (6)$$

Theorem

Suppose that $(\mathcal{C}, \mathcal{K})$ satisfies condition (6), and assume that for all $C \in \mathcal{C}$, $K \in \mathcal{K}$ with $C \prec K$ we have $|K \setminus C| = 1$. Then $(\mathcal{C}, \mathcal{K})$ is a proximity domain; pre-islands and islands corresponding to $(\mathcal{C}, \mathcal{K})$ coincide. Therefore, if $\mathcal{H} \subseteq \mathcal{C} \setminus \{\emptyset\}$ and $U \in \mathcal{H}$, then \mathcal{H} is a system of (pre-) islands if and only if \mathcal{H} is a distant family. Moreover, in this case \mathcal{H} is the system of (pre-) islands corresponding to its standard height function.

Example

Let $G = (U, E)$ be a connected simple graph with vertex set U and edge set E ; let \mathcal{K} consist of the connected subsets of U , and let $\mathcal{C} \subseteq \mathcal{K}$ such that $U \in \mathcal{C}$. Let \mathcal{C} consist of the connected convex sets of vertices.

Corollary

Let G be a graph with vertex set U ; let $(\mathcal{C}, \mathcal{K})$ be a connective island domain corresponding to $(\mathcal{C}, \mathcal{K})$, and let $\mathcal{H} \subseteq \mathcal{C} \setminus \{\emptyset\}$ with $U \in \mathcal{H}$. Then \mathcal{H} is a system of (pre-) islands if and only if \mathcal{H} is distant; moreover, in this case \mathcal{H} is the system of (pre-) islands corresponding to its standard height function.

THANK YOU FOR YOUR ATTENTION!

